

Datasheet



Mouse mAb to **MUC1 / EMA /
PEM / CD227**
Clone **VU-3C6**
Isotype **IgG1-κ**

Source

A BALB/c mouse was immunized with ZR-75-1 breast cancer cells.
Fusion partner: SP2/0.

Specifications

VU-3C6 reacts with the protein core of MUC1, an apical cell side epithelial marker which is upregulated or switched on in the majority of carcinomas. The dominant epitope of VU-3C6 is GVTSAPDTRPAP, located in the VNTR domain of MUC1. Binding of VU-3C6 is enhanced after glycosylation of the DTR motif. VU-3C6 was typed at the ISOBM TD-4 workshop.

Species reactivity

Positive: human.

Applications

VU-3C6 can be used for immunohistochemistry, ELISA and fluorescence tests.

ELISA	Frozen sections	Immunofluorescence	Paraffin sections
+	+	+	Periodic acid

Format

Produced in tissue culture, contains no host Ig. Antibodies are affinity purified and presented in PBS with 0,02 % sodium azide.

Stored at 4°C- 8°C, shelf life is at least 24 months after purchase.

Dilution advice

- ELISA (solid phase: 0,1-100 µg/ml; tracer: 0,001-100 µg/ml for 30 min at RT).
- Flow Cytometry (0,5-1,0 µg/million cells in 0,1 ml).
- Immunofluorescence (1-2 µg/ml).
- Immunohistology (formalin-fixed: 1-2 µg/ml for 30 min at RT; staining of formalin-fixed tissues is enhanced by preincubation of sections in 20mM periodic acid in 50mM acetate buffer pH 5.0 for 30 minutes at RT in the dark).

Positive control

MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.




Figure 1: Epitope of VU-3C6 as determined by epitope fingerprinting

Datasheet



References

- ISOBM TD-4 Workshop report, *Tumor Biol.* **19(Suppl 1)** (1998).
Karsten, U. et al. *Cancer. Res.* **58(12)**: 2541-2549 (1998).
Ryuko, K. et al. *Tumor Biol.* **21(4)**: 197-210 (2000).