

Datasheet



Mouse mAb to **Interferon α 1**
Clone **2-52**
Isotype **IgG1- κ**

Source

A BALB/c mouse was immunized with E. coli derived recombinant human IFN α 1.
Fusion partner: NS-0.

Specifications

The alpha interferons are involved in virus resistance in target cells for these viruses. They are known to block cell proliferation and to regulate MHC class I antigen expression. The IFN α family has over 20 genes and pseudogenes in two families (I and II), one with a mature length of 166aa and one of 172aa. Cells producing IFN α are lymphocytes, monocytes, macrophages and cell lines such as Namalwa and KGI. Bioassays for IFN α include cytopathic effect blocking, by viruses such as VSV, SFV and BMCV, on their target cells. A number of receptors for IFN α are now known and seem to be expressed on most cell types. 2-52 is specific for human IFN α 1 and does not cross react with human IFN α 2.



Figure 1:
Western blot
of HEK239T
stained for
IFN α 1

Species reactivity

Positive: human.

Applications

2-52 Can be used for the detection of human IFN α 1 in ELISA and Western blot. It can be paired with IFN α 1 mAb 2-48 to form an EIA to measure IFN α 1.

ELISA	Frozen sections	Pair	Western blot
+	+	2-48	+

Format

Produced in tissue culture, contains no host Ig. Antibodies are affinity purified and presented in PBS with 0,02 % sodium azide.

Stored at 4°C- 8°C, shelf life is at least 24 months after purchase.

Dilution advice

- ELISA (solid phase: 0,1-100 μ g/ml; tracer: 0,001-100 μ g/ml for 30 min at RT).
- Immunoblotting (1-2 μ g/ml).
- Immunohistology (1-2 μ g/ml for 30 min at RT; an appropriate antigen retrieval method for staining of formalin-fixed tissues has not been established to date).

Positive control

Human IFN α 1, Namalwa and KGI cells.

References

- Kontsek, P. et al., *Mol Immunol.* **29**: 863-870 (1992).